



Minireview

The clinical management of type 2 Gaucher disease

Karin Weiss¹, Ashley N. Gonzalez¹, Grisel Lopez, Leah Padoeim, Catherine Groden, Ellen Sidransky^{*}

Medical Genetics Branch, National Human Genome Research Institute, National Institutes of Health, Bethesda, MD, USA

ARTICLE INFO

Article history:

Received 23 September 2014

Received in revised form 6 November 2014

Accepted 7 November 2014

Available online 14 November 2014

Keywords:

Gaucher disease
Acute neuronopathic
Lysosomal storage disorder
Hydrops fetalis
Congenital ichthyosis
Palliative care

ABSTRACT

Gaucher disease, the inherited deficiency of the enzyme glucocerebrosidase, is the most common of the lysosomal storage disorders. Type 2 Gaucher disease, the most severe and progressive form, manifests either prenatally or in the first months of life, followed by death within the first years of life. The rarity of the many lysosomal storage disorders makes their diagnosis a challenge, especially in the newborn period when the focus is often on more prevalent illnesses. Thus, a heightened awareness of the presentation of these rare diseases is necessary to ensure their timely consideration. This review, designed to serve as a guide to physicians treating newborns and infants with Gaucher disease, discusses the presenting manifestations of Type 2 Gaucher disease, the diagnostic work-up, associated genotypes and suggestions for management. We also address the ethical concerns that may arise with this progressive and lethal disorder, since currently available treatments may prolong life, but do not impact the neurological manifestations of the disease.

Published by Elsevier Inc.

Contents

1. Introduction	111
2. Phenotypes associated with Type 2 GD	111
2.1. Hydrops fetalis	112
2.2. Skin abnormalities	112
2.3. Hepatosplenomegaly	112
2.4. Thrombocytopenia and anemia	112
2.5. Dysmorphology	112
2.6. Neurological involvement	112
3. Establishing the diagnosis of Gaucher disease	112
4. Genotype/phenotype correlation	112
5. Pathophysiology	113
6. The management of patients with Type 2 Gaucher disease	114
6.1. Failure to thrive and feeding difficulties	114
6.2. Bulbar involvement and airway compromise	114
6.3. Neurological aspects	118
6.4. Visceral and hematologic involvement	118
7. Treatment	119
7.1. Enzyme replacement and substrate reduction	119
7.2. Bone marrow and hematopoietic stem cell transplantation	119
8. Genetic counseling	119
9. Potential future therapies	119
9.1. Substrate reduction therapy	119
9.2. Chaperone therapy	119
9.3. Gene therapy	119
9.4. Additional treatment targets	119
10. Counseling parents and palliative care	119

^{*} Corresponding author at: Medical Genetics Branch, National Human Genome Research Institute, NIH, Room 1E623, 35 Convent Drive, Bethesda, MD 20892-3708, USA
E-mail address: sidranse@mail.nih.gov (E. Sidransky).

¹ These authors contributed equally.

11. Conclusion 120
 Acknowledgments 120
 References 120

1. Introduction

Gaucher disease (GD) is an autosomal recessive lysosomal storage disorder (LSD) caused by a deficiency of the enzyme glucocerebrosidase (GCase). It is the most common of the more than 50 known LSDs, and is traditionally divided into three distinct types based on the presence and progression of neurological signs [1]. Type 1, non-neuronopathic GD, presents with vast clinical heterogeneity, ranging from asymptomatic individuals to patients with severely enlarged organs and bone involvement [2,3]. Type 3 GD is considered the chronic neuronopathic form and is likewise accompanied by a spectrum of manifestations, often with visceral involvement as well as impaired horizontal saccadic eye movements, and some patients may develop myoclonic epilepsy, ataxia, or dementia [2,3]. Type 2 GD, or acute neuronopathic GD, presents with neurological involvement within the first months of life and progresses rapidly, culminating in severe neurodegeneration and death in infancy or early childhood [2,3].

While Type 2 GD is rare, its consideration is crucial for an accurate and timely diagnosis, appropriate management, genetic counseling and family planning [4]. It is encountered in all ethnicities. This work describes the signs and symptoms associated with Type 2 GD, management options, ethical concerns and how to establish the diagnosis. It is meant to serve as a guide for physicians and families of patients as they cope with a fatal diagnosis.

2. Phenotypes associated with Type 2 GD

Patients with Type 2 GD invariably exhibit clinical signs in the first year of life, although the type and severity of manifestations can vary widely [5,6]. While newborns often appear normal at birth and develop neurological signs with a rapid decline in health within the first year [6], others present prenatally or as neonates (Fig. 1). Although the visceral and neurological manifestations are fairly consistent, significant




Fig. 1. Establishing the diagnosis of Gaucher disease.

phenotypic heterogeneity is still observed. In the following sections we describe the key features reported.

2.1. Hydrops fetalis

Hydrops fetalis, characterized by an abnormal accumulation of fluid in multiple body areas, has been associated with several LSDs, including Type 2 GD [4]. While multiple disease processes are associated with hydrops, in over 25% of cases the cause is unknown. A portion of these undiagnosed cases may be due to LSDs that go unrecognized, leaving families unprepared regarding the risks for future pregnancies. Among patients with GD, cases involving hydrops tend to be perinatal lethal disorders and the pathophysiology is not well understood [6,7]. Generally, when hydrops is present in Type 2 GD, the fetus dies in utero, or is delivered prematurely and dies within several days of birth [7]. In cases of Type 2 GD without hydrops, fetuses generally reach full term [7].

2.2. Skin abnormalities

Newborns with Type 2 GD may present with congenital ichthyosis, an abnormal skin disorder resulting in shiny, almost cellophane-like skin. This is referred to as the collodion-baby phenotype [6,8]. The skin peels over the initial few weeks of life but then typically resolves while other symptoms worsen [6,9]. This pathology is thought to result from altered ratios of ceramides to glucosylceramides in the outermost layers of skin [10]. When severe, the skin must be treated aggressively with lubricants and fluids. Patients must be monitored vigilantly because of the risk for excessive loss of fluid and electrolyte abnormalities.

2.3. Hepatosplenomegaly

Hepatosplenomegaly, enlargement of both the liver and spleen, is one of the characteristic signs observed among patients with all three types of GD [7]. Often a first diagnostic clue can be the discovery of an enlarged spleen in the newborn nursery. Untreated, the enlargement can be massive, with organs that extend to the pelvic brim, compromising feeding and breathing. However, some infants may not exhibit prominent organomegaly.

2.4. Thrombocytopenia and anemia

Thrombocytopenia is observed among all forms of GD and occurs in approximately 40% of patients with Type 2 GD [7], either transiently during the neonatal period or as the disease progresses. Anemia is also frequently observed in these patients [6].

2.5. Dysmorphology

Dysmorphic facial features, often associated with other specific LSDs but generally not GD, have been described in some infants with Type 2 GD, including low-set ears and a small, upturned nose with a flat nasal bridge [6,7] (Fig. 2). Dysmorphic features often accompany the collodion-baby phenotype and might result from the presence of a thick membranous layer and decreased skin elasticity in utero.

2.6. Neurological involvement

All patients with Type 2 GD experience a rapid neurological decline, but manifestations vary widely. Some patients present with arthrogryphosis (joint contractures), microcephaly (abnormally small head circumference) or hypokinesia (decreased bodily movement) [6,7]. More common neurologic signs include a hyperextended neck, spontaneous apnea, hypertonica, and poor suck and swallow reflexes [6–8]. The characteristic facies and neck extension observed are thought to be secondary to a cervico-facial dystonia (Fig. 2). Strabismus

is also commonly seen. Many patients go on to develop seizures, rigidity and opisthotonos as the disease progresses.

3. Establishing the diagnosis of Gaucher disease

The recommended diagnostic workup for Type 2 GD is outlined in Fig. 3. The most reliable diagnostic methods are measurement of glucocerebrosidase enzymatic activity and direct sequencing of *GBA1*. Glucocerebrosidase enzymatic activity is usually included when ordering lysosomal panels, although it is important to assist the laboratory by providing appropriate clinical information. Also, unlike other lysosomal storage disorders, the enzymatic diagnosis of GD cannot be performed on plasma or RBCs, for only nucleated cells can be used. Enzymatic assays can be performed on leukocytes, fibroblasts, amniocytes or tissue samples such as liver or spleen. Type 2 GD is often difficult to distinguish from Types 1 and 3 based solely on enzymatic activity or genotype, although total absence of glucocerebrosidase is associated with early lethality.

Genetic testing for GD by direct sequencing of *GBA1* can be used to determine disease and carrier status [11]. Over 300 mutations in *GBA1* have been identified, and more continue to be discovered. Because there is a nearby pseudogene sharing 96% homology with *GBA1* in the coding regions, genotyping can be challenging as several mutations originate from the pseudogene sequence, and it is essential to differentiate between the two sequences. Further complicating molecular diagnosis, many mutations are rare or private point mutations, insertions, deletions or splice-site changes that may not be detected in limited mutation screens. Some mutations appear to have arisen as a result of recombination event that occurred between *GBA1* and its homologous nearby pseudogene and are referred to as recombinant alleles. Recombinant alleles are relatively common among infants with Type 2 GD. Since recombinant alleles may result in the introduction of more than one alteration on the same allele, the entire sequence must be carefully analyzed. Generally, the enzyme assay is not reliable for heterozygote detection and DNA tends to be far more accurate, especially in families where the *GBA1* mutations are known.

Patients with Type 2 GD exhibit epidermal abnormalities regardless of whether ichthyosis is clinically evident [9,12]. This finding may have potential diagnostic importance. On electron microscopy, the skin ultrastructure reveals immature partially-processed lamellar bilayers. A study comparing 20 cases with Type 2 GD to other forms of GD demonstrated that these alterations were unique to Type 2 GD and could help differentiate between patients with Type 2 versus Type 3 GD. However, such evaluations are currently performed on a research-only basis [13].

While it is not a recommended method for diagnosis, many affected infants have been identified as a result of pathologic evaluations of biopsy specimen. The classic finding in bone marrow biopsies or liver, lung, and spleen samples is the presence of Gaucher cells, lipid engorged macrophages with a “wrinkled tissue” appearance of the cytoplasm. However, enzymatic or DNA confirmation is essential, as storage cells resembling Gaucher cells can be found in other conditions.

Clinicians should consider a diagnosis of GD in newborns with hepatosplenomegaly, hematological abnormalities, developmental delay, and failure to thrive. Among Ashkenazi Jewish families the carrier frequency is about 1 in 16; however neuronopathic forms of GD are relatively uncommon in this population [14].

4. Genotype/phenotype correlation

It is generally difficult to predict disease progression in GD based on the *GBA1* genotype. Table 1 summarizes the genotypes and clinical features of 77 patients with Type 2 GD described in the literature. The nomenclature for each amino acid change encountered in *GBA1* has recently been revised to take into consideration the 39 amino acid leader sequence and thus mutation reports can be a source of some confusion. We have included both designations with the newer versions in



Fig. 2. Two patients with Type 2 Gaucher disease demonstrating the characteristic faces and strabismus. Top panel shows the first infant at age 6 months (left) and post-tracheotomy at 16 months (right). Lower panel shows the second child at age 8 months (left) and after tracheotomy at age 2 years.

parentheses. It is evident that there are numerous genotypes and clinical phenotypes associated with Type 2 GD. However, certain generalizations can be made based on these data, notably the absence of one of the most common *GBA1* mutations, Asn370Ser (Asn409Ser) [15]. Patients heterozygous or homozygous for this mutation do not manifest Type 2 GD. There has, however, been at least one report where Asn370Ser (Asn409Ser) was found on the same allele in cis with another *GBA1* mutation, which can further confuse the picture [16]. In contrast, the mutation Leu444Pro (Leu483Pro) is frequently encountered in Type 2 GD, although it is rare to identify patients who are homozygous. When homozygosity for this mutation is reported in patients with Type 2 GD, further evaluation often shows that at least one Leu444Pro (Leu483Pro) allele is part of a recombinant allele with other mutations. Such alleles are null alleles, and homozygosity results in early lethality. Other null alleles like c.84insG or IVS2+1G>A are also found in Type 2 GD, often with a Leu444Pro (Leu483Pro) mutation on the second allele. Mutations involving an arginine such as Arg257Gly (Arg286Gly), Arg285His (Arg324His), Arg131Leu (Arg170Leu), Arg120Try (Arg159Try), Arg359X (Arg398X), and Arg463His (Arg463His) appear to be frequent. One allele with two *GBA1* mutations, D409H + H255Q (Asp448His + His294Gln), is often found among patients with Type 2 GD of Greek or Albanian ancestry [17]. A relatively frequent *GBA1* polymorphism E326K (E365K) is found in roughly 2–4% of the general population and can also be encountered among patients with Type 2 GD along with a second *GBA1* mutation on the same allele.

5. Pathophysiology

It is still not known why only a subset of patients with GD develops aggressive neurodegeneration. Elucidating the etiology has additional

relevance because mutations in *GBA1* are a risk factor for developing Parkinson disease [18]. The deficiency of GCCase in patients leads to the accumulation of glucosylceramide (GlcCer) and glucosylsphingosine (GlcSph) in many tissues, including the brain. Measurements of GlcCer and GlcSph in brain samples from patients with GD reveal that GlcSph is greatly elevated in neuronopathic GD, with the highest levels seen in Type 2 [19]. Recent work by Futerman and colleagues using a mouse model of neuronopathic GD found that the continued accumulation of GlcCer reaches a critical threshold that triggers a rapid cascade leading to neuroinflammation and neurodegeneration in certain brain regions [20,21].

The role of necroptosis in the pathogenesis of Type 2 GD has also been explored and considered as a potential therapeutic target based on work in animal models [22]. Cell death occurs in a regulated manner via necrotic mechanisms, beginning with the formation of the necrosome, a multiprotein complex which includes receptor-interacting protein kinase-1 (RIPK1) and RIPK3 [23]. Necrosome formation can be initiated by ligand binding to death receptors and toll-like receptors, among others [23]. A caspase inhibitor must be present to prevent the complex from following a caspase-dependent apoptotic pathway [23]. Modification of RIPK1 and RIPK3 activate the kinases, beginning a downstream signaling cascade resulting in necroptotic cell death [23]. Active RIPK1 and RIPK3 levels were elevated in brain tissue of Gaucher mice generated by a conditional knock-out strategy. Furthermore, *Ripk3*^{-/-} knock-out mice treated with the highly specific glucocerebrosidase inhibitor, conduritol-beta-epoxide, had fewer neurologic and systemic manifestations and greater longevity compared to similarly treated control mice [22]. These observations suggest a role for necroptosis in neuronopathic forms of GD, but further studies are needed in human tissues.



Fig. 3. Neonatal presentations of Gaucher disease demonstrating organomegaly (top left), hydrops (bottom) and peeling skin (top right).

6. The management of patients with Type 2 Gaucher disease

Management of Type 2 GD can be challenging for physicians as well as for the patients' families. It is not the intent of this article to present rigid management guidelines, as many cultural, financial and ethical considerations come into play. Common management challenges and suggestions are also outlined in Table 2.

6.1. Failure to thrive and feeding difficulties

Failure to thrive is a common finding in patients with Type 2 GD and results from a combination of anorexia, nausea, hypotonia or hypertonia, vomiting and feeding difficulties [6]. Chewing difficulties, a weak suck, and/or problems with swallowing occur due to neurological involvement. In addition, the presence of abnormal neck tone and opisthotonos, trismus and/or the presence of laryngospasm can be contributing factors [24]. When the diagnosis is reached at an early stage, oral feeding may still be possible. A clinical evaluation of feeding should be performed by a speech pathologist, and infants should be monitored for weight gain, dehydration and/or signs of aspiration [25]. A video fluoroscopic evaluation of swallowing (modified barium scan) provides further information regarding the presence of aspiration, gastro-esophageal reflux and laryngeal penetration with different types of feedings [25] and can be used to adjust the type of bottle/nipple, formula thickening or recommend strict enteral feeding. It can also provide objective evidence for the extent of bulbar involvement [24,26]. The images can be used as an educational tool in discussions regarding the advantages and disadvantages of oral versus gavage feedings. In cases when it is decided to switch to enteral feedings, a nasogastric tube or gastric tube insertion with gastrostomy are two acceptable approaches. Decisions regarding the most appropriate approach should take into account potential anesthesia-related complications during a gastrostomy or gastric tube insertion procedure [27].

Assessment of macronutrient, micronutrient and fluid intake is important in these patients because their oral intake is usually low. An individualized nutrition plan should be constructed, considering the mode of feeding and energy requirements. Patients with Type 2 GD may not gain weight even with enteral or parenteral feeding because of increased resting energy expenditure [28].

6.2. Bulbar involvement and airway compromise

Brain stem degeneration leads to bulbar involvement, characterized by choking events, laryngeal spasm and apnea. These frightening episodes occur in a subset of patients and tend to occur with increased frequency as the disease progresses [29]. Laryngospasm can also lead to hypoxemia, which may accelerate the disease process. Treatment of gastro-esophageal reflux and strict enteral feeding can decrease the incidence of these events in early stages of the disease, but spontaneous episodes may occur as the disease worsens [29]. Excessive oral secretions may also contribute to respiratory symptoms, necessitating suction. Treatment with sublingually delivered ophthalmic atropine drops has been used with some success [30].

Whether to place a tracheotomy tube is one of the most challenging issues faced by physicians and families caring for children with Type 2 GD. A tracheostomy has been effectively used when the primary manifestation is laryngospasm and airway compromise. While in general patients with Type 2 GD who undergo tracheostomy placement tend to survive longer, consideration that this intervention may prolong suffering should be discussed with the family [31]. Extensive dialogue regarding realistic expectations is desirable prior to undertaking this measure. Parents should be counseled that the decision not to undergo tracheotomy can be very reasonable in many circumstances.

There is an increased risk for anesthesia-related complications due to increased secretions, gastro-esophageal reflux, poor control of laryngeal musculature and/or laryngeal spasm. These may be triggered by

Table 1
Genotypes and significant clinical features observed in patients with Type 2 Gaucher disease.

Alleles	Protein	cDNA	Age at diagnosis	Age at death	Hydrops	Ichthyosis	Hepatosplenomegaly	Seizures	Additional clinical information	Reference
c.1263del55/Met416Val	p.Met455Val	c.1263del55/ c.1363A>G	8 months	17 months				Yes		[12]
c.1263del55/Arg285His	p.Arg324His	c.1263del55/ c.971G>A	At birth	Perinatal					Unknown	[13]
c.1263del55/Arg257Gln	p.Arg296Gln	c.1263del55/ c.887G>A	At birth	3 weeks		Yes		Yes		[13]
Ala190Glu/Gly325Arg	p.Ala229Glu p.Gly364Arg	c.686C>A/ c.1090G>A	9 months	Unknown					Unknown	[13]
Arg120Trp/Ser196Pro	p.Arg159Trp p.Ser235Pro	c.475C>T/c.703T>C	At birth	3 months		Yes		Yes Yes	Laryngospasm	[70]
Arg120Trp/Ser196Pro	p.Arg159Trp p.Ser235Pro	c.475C>T/c.703T>C	At birth	11 days		Yes		Yes Yes (refractory)		[13]
Arg131Leu/Arg131Leu	p.Arg170Leu	c.509G>T/c.509G>T	5 months	7 months		Yes			Developmental regression	[13]
Arg131Leu/Arg131Leu	p.Arg170Leu	c.509G>T/c.509G>T	Prenatal	9 months		Yes		Yes		[71]
Arg359X/Val398Phe	p.Arg398X p.Val437Phe	c.1192C>T/ c.1309G>T	At birth	4 weeks		Yes			Facial dysmorphism (triangular face, malformed ears)	[72]
Arg463His/Arg463His	Arg502His Arg502His	c.1505G>A/ c.1505G>A	At birth	1 month		Yes		Yes Yes (myoclonic)	Respiratory failure	[73]
Asp409His+His255Gln/ Asp409His+His255Gln	Arg448His+ His294Gln/ Asp448His+ His294Gln	c.[1342G>C c.882T>G]/ c.[1342G>C c.882T>G]	1 month	Unknown					Unknown	[74]
Asp409His+His255Gln/ Asp409His+His255Gln	Asp448His+ His294Gln/ Asp448His+ His294Gln	c.[1342G>C c.882T>G]/ c.[1342G>C c.882T>G]	8 months	2 years					Brainstem involvement; developmental regression	[74]
c.533delC/c.533delC		c.533delC/ c.533delC	Prenatal	22 weeks gestation	Yes			Yes	External abnormalities (rocker bottom feet, flat nose)	[5,75]
c.84dupG/Asp409His	Asp448His	c.84dupG/ c.1342G>C	5 months	Unknown					Abnormal eye saccades; delayed cognition	[74]
Cys16Ser/Cys16Ser	Cys55Ser/Cys55Ser	c.164G>C c.164G>C	Postmortem	At birth					Unknown	[76]
Cys16Ser/Cys16Ser	Cys55Ser/Cys55Ser	c.164G>C c.164G>C	Prenatal	33 weeks gestation					4th unsuccessful pregnancy for couple	[76]
Gly202Arg/Gly202Arg	Gly241Arg/ Gly241Arg	c.721G>A c.721G>A	5 months	Unknown					Unknown	[13]
Gly202Arg/Gly202Arg	Gly241Arg/ Gly241Arg	c.721G>A c.721G>A	6 months	Unknown					Unknown	[13]
His311Arg/His311Arg	His350Arg/ His350Arg	c.1049A>G c.1049A>G	Prenatal	At birth	Yes			Yes		[72]
His311Arg/His311Arg	His350Arg/ His350Arg	c.1049A>G c.1049A>G	Prenatal	At birth	Yes			Yes		[72]
IVS2+1G>A/Phe251Leu	Phe290Leu	c.115+1G>A/ c.870C>A	16 days	1 month					Decreased spontaneous movement	[74]
IVS2+1G>A/Phe251Leu	Phe290Leu	c.115+1G>A/ c.870C>A	At birth	1 month		Yes		Yes Myoclonic jerks/ tremors		[77]
IVS2+1G>A/RecNcil		c.115+1G>A ^a	Prenatal	45 days	Yes	Yes		Yes		[78]
IVS2+1G>A/RecNcil		c.115+1G>A ^a	Unknown	Unknown					Oculomotor apraxia; developmental regression	[74]
Leu444Pro+Glu326Lys/ Pro182Leu	Leu483Pro+ Glu365Lys/ Pro221Leu	c.[1448T>C; c.1093G>A]/ c.662C>T	At birth	25 days	Yes			Yes		[79]

(continued on next page)

Table 1 (continued)

Alleles	Protein	cDNA	Age at diagnosis	Age at death	Hydrops	Ichthyosis	Hepatosplenomegaly	Seizures	Additional clinical information	Reference
Leu444Pro/Arg131Cys	Leu483Pro/Arg179Cys	c.1448T>C/ c.508C>T	8 months	10.5 months				Yes	Strabismus	[80]
Leu444Pro/Arg257Gln	Leu483Pro/Arg296Gln	c.1448T>C/ c.887G>A	Unknown	Unknown					Extensor posturing of trunk and neck; dysconjugate and crossed gaze	[74]
Leu444Pro/Arg257Gln	Leu483Pro/Arg296Gln	c.1448T>C/ c.887G>A	9 months	3 years				Yes (myoclonic)		[81]
Leu444Pro/Arg257Gln	Leu483Pro/Arg296Gln	c.1448T>C/ c.887G>A	7 months	Unknown				Yes	Strabismus; bilateral hearing loss; failure to thrive	[77]
Leu444Pro/Asp380Asn	Leu483Pro/Asp419Asn	c.1448T>C/ c.1255G>A	Unknown	Unknown					Unknown	[74]
Leu444Pro/c.330delA	Leu483Pro	c.1448T>C/ c.330delA	3 months	Unknown					Unknown	[13]
Leu444Pro/Gln414X	Leu483Pro/Gln453X	c.1448T>C/ c.1357C>T	9 months	10 months					Apneic spells; developmental regression; ophthalmoplegia	[74]
Leu444Pro/Gln414X	Leu483Pro/Gln453X	c.1448T>C/ c.1357C>T	2 months	10 months					Unknown	[74]
Leu444Pro/Gly202Arg	Leu483Pro/Gly241Arg	c.1448T>C/ c.721G>A	8 months	2.5 years				Yes (tonic)		[81]
Leu444Pro/Ile260Thr	Leu483Pro/Ile299Thr	c.1448T>C/ c.896T>C	6 months	21 months					Strabismus; facial dysmorphism; respiratory difficulties	[79]
Leu444Pro/IVS2+1G>T	Leu483Pro	c.1448T>C/ c.115+1G>T	6 months	9 months					Brainstem involvement; developmental regression	[74]
Leu444Pro/IVS2+1G>A	Leu483Pro	c.1448T>C/ c.115+1G>A	4 months	Unknown					Unknown	[13]
Leu444Pro/IVS2+1G>A	Leu483Pro	c.1448T>C/ c.115+1G>A	4 months	Unknown					Unknown	[13]
Leu444Pro/IVS2+1G>A	Leu483Pro	c.1448T>C/ c.115+1G>A	6.5 months	1 year					Strabismus; hypotonia; swallowing difficulties	[82]
Leu444Pro/IVS2+1G>A	Leu483Pro	c.1448T>C/ c.115+1G>A	At birth	9 months				Yes	Progressive difficulty swallowing; bilateral brain stem dysfunction	[82]
Leu444Pro/IVS2+1G>T	Leu483Pro	c.1448T>C/ c.115+1G>T	Unknown	8.5 months					Apnea; hypotonia; hyperflexia; esotropia	[74]
Leu444Pro/Leu385Pro	Leu483Pro/Leu424Pro	c.1448T>C/ c.1271T>C	11 months	18 months				Yes	Ophthalmoplegia	[12]
Leu444Pro/Leu444Pro+Ala456Pro	Leu483Pro/Leu483Pro+Ala495Pro	c.1448T>C/ c.[1448T>C c.1453G>C]	7 months	10 months			Yes		Strabismus	[82]
Leu444Pro/Phe213Ile	Leu483Pro/Phe252Ile	c.1448T>C/ c.754T>A	1 year	28 months					Ophthalmoplegia	[12]
Leu444Pro/Phe213Ile	Leu483Pro/Phe252Ile	c.1448T>C/ c.754T>A	11 months	Unknown					Bone involvement; ophthalmoplegia	[12]
Leu444Pro/RecNcil	Leu483Pro	c.1448T>C ^d	5 months	1 year				Yes	Strabismus; facial paralysis; poor gag	[8]
Leu444Pro/RecNcil	Leu483Pro	c.1448T>C ^d	1 year	22 months					Abnormal tone	[12]
Leu444Pro/RecNcil	Leu483Pro	c.1448T>C ^d	8 months	18 months				Yes	Ophthalmoplegia	[12]
Leu444Pro/RecNcil	Leu483Pro	c.1448T>C ^d	1 month	6 months					Unknown	[74]
Leu444Pro/Thr323Ile	Leu483Pro/Thr362Ile	c.1448T>C/ c.1085C>T	6 months	Unknown				Hand tremors	Strabismus; brainstem involvement; clonus in feet	[74]
Leu444Pro/Tyr304Cys	Leu483Pro/Tyr343Cys	c.1448T>C/ c.1028A>G	1 year	Unknown					Unknown	[13]
Leu444Pro/Tyr304Cys	Leu483Pro/Tyr343Cys	c.1448T>C/ c.1028A>G	1 year	2 years				Yes (myoclonic)		[81]
Leu444Pro/Tyr304Cys	Leu483Pro/Tyr343Cys	c.1448T>C/ c.1028A>G	1 year	Unknown					Unknown	[74]
Leu444Pro/Tyr313His	Leu483Pro/Tyr352His	c.1448T>C/ c.1054T>C	7 months	10 months					Oculomotor apraxia; developmental regression	[74]
Leu444Pro/unknown	Leu483Pro	c.1448T>C	7 months	11 months				Yes Yes	Strabismus	[8]

Leu444Pro/unknown	Leu483Pro	c.1448T>C	Unknown	Unknown				Unknown	[74]
Leu444Pro/unknown	Leu483Pro	c.1448T>C	8 months	2.5 years	Yes	Yes		Absent gag	[8]
Lys198Glu/Lys198Glu	Lys237Glu/ Lys237Glu	c.709A>G/ c.709A>G	16 months	2 years	Yes	Yes (myoclonic)			[81]
Lys198Glu/Lys198Glu	Lys237Glu/ Lys237Glu	c.709A>G/ c.709A>G	Unknown	Unknown			Myoclonic jerks	Hypotonia; gaze abnormalities	[74]
Phe213Ile/Asn382Lys	Phe252Ile/ Asn421Lys	c.754T>A/ c.1263C>A	10 months	18 months				Unknown	[12]
Phe213Ile/Leu383Arg	Phe252Ile/ Leu422Arg	c.754T>A/ c.1265T>G	7 months	26 months			Yes		[12]
Phe213Ile/Lys74X	Phe252Ile/Lys113X	c.754T>A/c.337A>T	Unknown	10 months			Yes (myoclonic)	Strabismus; ophthalmoplegia; apneic spells	[74]
Phe259Leu/Asn188Lys	Phe298Leu/ Asn227Lys	c.894C>A/c.681T>G	At birth	Perinatal				Opisthotonus	[13]
Pro122Leu/Val375Leu	Pro161Leu/ Val414Leu	c.482C>T/ c.1240G>C	1 year	Unknown			Yes		[12]
RecA/IVS 10+2		c.1505+2T>G ^a	Prenatal	Prenatal	Yes	Yes		Ascites; pleural effusions	[74]
RecA/IVS 10+2T>G		c.1505+2T>G	Prenatal	Prenatal termination	Yes				[82]
RecB/RecB		^a	Prenatal	1 day	Yes	Yes	Yes	Facial dysmorphism	[8]
RecB/RecB		^a	Prenatal	Miscarriage	Yes			Thymic hypoplasia	[8]
RecD/RecD		^a	Prenatal	Perinatal	Yes				[13]
RecD/RecD		^a	Prenatal	2 hours	Yes	Yes	Yes		[83]
RecD/RecD		^a	Prenatal	1 hour	Yes	Yes	Yes	Hypokinesia; facial dysmorphism	[84]
RecG+Leu444Pro/ Arg257Gln	Leu483Pro/ Arg296Gln	c.1448T>C/ c.887G>A	6 months	Unknown				Unknown	[13]
RecNcil/Glu41Leu	Glu80Leu	c.238G>A ^a	Several months	18 months			Yes	Failure to thrive	[80]
RecNcil/RecNcil		^a	Prenatal	At birth	Yes		Yes	Hyperextension of neck	[71]
Ser196Pro/Ser196Pro	Ser235Pro/ Ser235Pro	c.703T>C/c.703T>C	At birth	Perinatal				Respiratory difficulty	[13]
Trp184Arg/Asp409His	Trp223Arg/ Asp448His	c.667T>C/c.1342G>C	5 months	Unknown				Opisthotonos; hypertonia; increased gag reflex	[74]
Tyr363Cys/unknown	Tyr402Cys	c.1205A>G	5 months	15 months			Yes		[12]
Unknown/unknown			At birth	2 months			Yes	Hyperextension of the neck; hypertonia; poor suck/swallow	[85]
Unknown/unknown			At birth	1 week		Yes		Flat nose	[85]

^a Rec D: recombination at the end of exon 7; Rec G: recombination in intron 10-exon 11 or the 3' untranslated region; Rec A: Recombination in intron 3; Rec Ncil: Recombinant allele in exon 10 containing L444P, A456P, and V460V.

Table 2
Managing common clinical features of Type 2 GD.

Clinical feature	Recommended evaluation	Management
Feeding difficulties and failure to thrive	Exam by a speech pathologist Video fluoroscopy to assess swallowing and aspirations Nutrition assessment	Modification of bottle type/formula thickening Consider strict enteral feeding per nasogastric tube/gastric tube Individualized nutrition plan with sufficient macro and micronutrients
Gastrointestinal reflux	Clinical diagnosis Can be demonstrated on video fluoroscopy and pH meter	Head elevation Proton pump inhibitors/H2 blockers
Laryngeal spasm/choking/apnea	Consider ENT evaluation	Treat GE reflux Atropine drops for excess salivation Consider strict enteral feeding Consider tracheostomy although may prolong survival Take precautions when performing anesthesia Occupational therapy evaluation
Spasticity/abnormal movements	Neurologic evaluation	Consider benzodiazepines with careful monitoring Consider EEG
Irritability	Nutrition assessment of intake	Provide sufficient macro- and micronutrients Gentle handling and maintenance of a calm and familiar environment Consider benzodiazepines with careful monitoring
Fever	Evaluate for infection/recurrent aspirations	Antipyretics Consider empiric antibiotic treatment
Lung disease	Monitor for oxygen desaturations/respiratory distress on physical exam Chest X ray for evaluation of disease progression and pulmonary infections Video fluoroscopy for evaluation of aspirations Consider pulmonology evaluation	Treat GE reflux Treat pulmonary infections Oxygen
Visceromegaly	Monitor CBC	Enzyme replacement therapy is generally not indicated
Bleeding diathesis	Check CBC, PT/PTT Check liver function tests Consider testing coagulation factors	Vitamin K Consider transfusion of platelets/fresh frozen plasma

pre-anesthesia medications or manipulation of the airway [27]. Difficult airway access may require the need for smaller endotracheal tubes [27]. Pulmonary involvement in GD is another risk factor for anesthesia-related complications.

6.3. Neurological aspects

Neurological involvement can manifest as irritability and spasticity. Non-pharmacological treatments include gentle handling, maintaining a calm and familiar surrounding, physical therapy and proper positioning [32]. Pharmacological treatment with benzodiazepines can be useful for both irritability and spasticity. Caregivers should be educated about possible side effects including sedation, constipation, urinary retention, hypersalivation and weakness. Respiratory depression can be seen with benzodiazepine toxicity. Airway compromise due to bulbar involvement may limit their use. Other anti-spasticity drugs used include baclofen, dantrolene and tizanidine [33]. These medications can be offered in the setting of palliative therapy [32].

Seizures, sudden episodes of sensory disturbance, loss of consciousness, or convulsions, associated with abnormal electrical activity in the brain, can occur as the disease progresses. Electroencephalography (EEG) can identify abnormal brain activity in the absence of clinical signs. Findings on EEG may include slowing and disorganization of the background and/or epileptiform activity [28,33]. Myoclonic epilepsy and myoclonus are often described although not universally, and are thought to be mainly of cortical origin [29,34].

Seizure control can be very challenging in patients with Type 2 GD. Careful attention to the identification of treatable etiological factors such as hypoglycemia, hypocalcemia, hyponatremia, hypomagnesemia, hypoxia, intracranial hemorrhage, hyperammonemia, and/or infections is of critical importance. High quality neuroimaging and accurate EEG interpretation may help in optimizing patient care and outcomes [35]. Continuous EEG monitoring for the detection of subclinical seizures is recommended, but may not be possible due to the need for specialized training and access to equipment [36]. Periodic conventional EEG monitoring is more generally available. Antiepileptic treatment is generally

guided by the particular epilepsy syndrome at presentation [37]. However, benzodiazepines are usually first-line therapy for status epilepticus although they can cause respiratory depression and/or cardiac dysrhythmia. Midazolam and lorazepam are the preferred agents. Second-line therapies include phenytoin and fosphenytoin. Phenytoin has been associated with local skin necrosis and hypotension. Both agents can cause cardiac dysrhythmias with more rapid administration. Phenobarbital is frequently used in infants, but carries the risk of hypotension, respiratory depression, and cardiac arrhythmia. Levetiracetam can also be considered. Valproate can be effective but patients must be monitored for potential side effects such as thrombocytopenia and hepatotoxicity [38].

6.4. Visceral and hematologic involvement

Visceral involvement presents as splenomegaly and is associated with anemia and thrombocytopenia. Liver enlargement is also seen, and can be accompanied by elevations in transaminases. Portal hypertension and liver dysfunction are rarely described in Type 2 GD, although neonatal cholestasis is occasionally seen and may precede other manifestations [39,40]. Splenectomies have been performed in these patients but are generally not recommended due to the risk of surgical complications. It is important to note that the absence of overt organomegaly is not sufficient to rule out the diagnosis of Type 2 GD.

The main hematological complication is a bleeding tendency secondary to the thrombocytopenia and abnormal platelet aggregation [41]. A decrease in clotting factors secondary to consumption by an enlarged spleen has also been described [42]. The complete blood cell count should be monitored regularly to identify the need for transfusions. Systemic disease can be minor or almost absent in comparison to the neurological deficits observed in patients with Type 2 GD.

Pulmonary involvement with interstitial lung disease is a common finding primarily resulting from recurrent aspirations [6] or from the infiltration of alveolar, interstitial, perivascular, and peribronchial spaces by lipid-laden macrophages [43]. Chest X-ray may show diffuse interstitial infiltration in both lung fields, and a ground glass pattern can be seen on

chest CT. On lung biopsy, Gaucher cells can be seen [43]. Periodic monitoring of pulmonary involvement should include examination for evidence of respiratory distress, oxygen saturation measurements, and chest X-rays. Such testing can identify the need for supportive treatment with oxygen and/or antibiotics if there is evidence for aspiration pneumonia. Fever may be a sign of a pulmonary infection, but can also be present in the absence of infection due to inflammation, possibly mediated by activated glucosylceramide-laden macrophages [6]. Prolonged, unexplained fever has been associated with a worsening disease course [29].

7. Treatment

7.1. Enzyme replacement and substrate reduction

Enzyme replacement therapy (ERT) can ameliorate the visceromegaly and hematologic abnormalities in patients with Type 2 GD [28,44]. It is not clear whether ERT impacts pulmonary involvement, although when the primary cause of pulmonary involvement is recurrent aspirations secondary to neurological deterioration, ERT is not beneficial [28,44]. The recombinant enzyme does not cross the blood–brain barrier and there is no evidence that ERT has reversed, stabilized, or slowed the progression of neurological involvement [45]. Siblings were described where one received ERT after diagnosis at 7 months, and the subsequent sibling was treated from birth. While the sibling treated pre-symptomatically survived 6 months longer, the final outcome was the same [45]. It has been debated whether it is ethical to treat Type 2 GD patients with ERT because it is a very costly and inconvenient therapy that may only prolong suffering [46]. It is generally accepted that ERT should be discussed with families, presenting both its potential merits and limitations. Some physicians feel that ERT is not indicated for patients with Type 2 GD, although in recent years many infants have been started on the therapy [45]. It is sometimes deemed appropriate to treat with ERT until it is clear that the patient does not have Type 3 GD. In addition, it is reasonable to consider therapy for Type 2 GD patients in situations where it could be palliative, for example if reduction of organomegaly would alleviate pain, avoid surgery or facilitate other necessary interventions such as gastric tube placement.

Alternative methods of enzyme delivery have been investigated to treat the neurological deterioration in GD. Intraventricular delivery via a reservoir was attempted without success [47]. An animal study evaluated convection-enhanced delivery (CED) to restore GCase levels in the brains of rats and primates [48]. Enzymatic assays showed significant increases in GCase activity in the white matter, cortex, and pons of treated animals, and immunohistochemical staining confirmed GCase localization in neurons of the cortex [48]. One patient with Type 2 GD was treated with GCase via CED, and while successful delivery of enzyme was documented, there was continued disease progression [49]. Intravascular delivery after disruption of the blood–brain barrier has been investigated but not attempted in these patients [50].

Substrate reduction therapy with Miglustat, an inhibitor of glucosylceramide synthase, was not shown to be beneficial for neurological manifestations in Type 3 GD [51]. This treatment is not indicated for Type 2 GD.

7.2. Bone marrow and hematopoietic stem cell transplantation

Full engraftment of a hematopoietic stem cell transplantation (HSCT) has been performed in patients with Type 1 GD resulting in complete hematologic correction [52,53]. Ringden et al. described four Swedish patients with Type 3 GD who underwent bone marrow transplantation between the ages of 2–9 years old [54]. Two patients had no cognitive decline at follow-up ten years later [54]. However, both developed epilepsy 14 and 22 years after allo-HSCT [55]. Thus, the procedure does not prevent development of neurological damage. Published reports regarding the results of bone marrow transplantation in

patients with Type 2GD are not available. The risks involved in transplanting a seriously ill infant must be seriously considered, given the strong possibility of continued neurological deterioration. Moreover, end-of-life issues may need to be addressed in distant centers, away from the support of home and family.

8. Genetic counseling

Genetic counseling is an important aspect of the care for these patients and families. Parental feelings of guilt or blame can complicate the decision-making process as the child's condition deteriorates. Families need to be educated about risks and alternatives relevant to future family planning, including prenatal diagnosis, pre-implantation diagnosis, sperm or egg donation, as well as adoption. In addition, the genetics team should be aware that unusual mechanisms including uniparental disomy for chromosome 1 and germline or de novo mutations have been described in patients with GD [56,57]. Thus the recurrence risk is not always obvious.

9. Potential future therapies

9.1. Substrate reduction therapy

Inhibitors of glucosylceramide synthase have been explored as a means to attenuate glucosylceramide production and accumulation in various organs. The currently available therapy is not beneficial in neuropathic GD (Type 2 and 3) [51]. A novel glucosylceramide synthase inhibitor decreased substrate accumulation in the brains of a mouse model for neuronopathic GD [58]. This treatment modality might be beneficial for neuronopathic GD if given pre-symptomatically and before significant accumulation of substrate in the brain occurs, perhaps through newborn screening programs [59].

9.2. Chaperone therapy

Restoration of the mutant enzyme activity by small molecule compounds (chaperones) capable of crossing the blood–brain barrier is a new approach to treat LSDs [60,61]. Currently, there is an effort to identify candidate molecules [62,63]. However, this treatment will be helpful only for mutations in *GBA1* that result in an unstable mutant protein and may not be appropriate for patients with null alleles. Induced pluripotent stem cells (iPSCs) generated from fibroblasts of patients with Type 2 GD may facilitate studies of candidate molecules. Gaucher iPSC lines show deficient GCase and have been differentiated into macrophages that exhibit substrate accumulation [64]. This GD phenotype in macrophages was successfully reversed with a new molecular chaperone [65].

9.3. Gene therapy

Gene therapy is a potentially promising future therapeutic strategy for genetic diseases with enzymatic deficiencies like GD [66], although the effectiveness of gene therapy in CNS disease is not clear. Intracerebral injection of the vector carrying the wild-type gene could be a possible strategy [66].

9.4. Additional treatment targets

As we advance our understanding of the mechanisms involved in neurodegeneration, other potential targets will likely be identified. Insights from basic science research such as the potential role RIPK3 [22] may provide new leads for future drug development.

10. Counseling parents and palliative care

Facing the diagnosis of Type 2 GD in an infant is extremely difficult and can be emotionally overwhelming for parents, caregivers and

health care providers. In many cases, the child appears normal at birth and during the first months of life until rapid deterioration begins. In the midst of these circumstances, parents and clinicians should work together to make important and complex decisions about the child's medical care. These decisions are based on a variety of considerations including medical information, values, cultural expectations, economic considerations and parental beliefs [67]. Studies have shown that parents of critically ill children need sensitivity, empathy, follow-up contact and information counseling from healthcare providers [68]. This combination of education, support, and empathy can best be achieved by a multidisciplinary approach. Patient support groups or on-line chat rooms can also be helpful to some parents.

Type 2 GD is a progressive condition with an aggressive neurodegenerative course and is therefore particularly appropriate for palliative care. This can be a supportive resource for the parents as well as the child. When it is evident that supportive measures are the only options appropriate for continued care, health care providers can help the parents reengage with a new set of goals. The new goals can include managing the child's condition with the least amount of treatment-related pain or suffering, limiting exposure to invasive or extreme interventions, and maintaining the child's quality of life [67]. This approach might seem very reasonable for some parents and not to others, who choose to continue intensive interventions. Cultural and religious beliefs may affect the parental decisions on palliative and end of life care and must be considered [68]. It is important to provide the parents with ongoing counseling by healthcare professionals during the different stages of the disease as well as after the child's death [69].

When possible, the subject of an autopsy should be raised with families. Such post-mortem evaluations can help clinicians and researchers to better understand disease pathogenesis. Frozen stored tissue specimens are very useful for many studies that could ultimately impact the care of future children afflicted with this disorder.

Caring for a dying child can have a great impact on both parents and siblings. Appropriate support and therapy should be provided as needed. Also, with the parents' consent, education of employers, extended family members, and clergy as to the gravity of the illness can help the family get the needed support and time with their child. Engaging the services of hospice can also be very helpful.

11. Conclusion

Diagnosing and treating infants with Type 2 Gaucher disease represents a challenging paradigm for neonatal and genetic illnesses. An accurate and timely diagnosis of GD is critical for genetic counseling, patient management and family planning that can inform future pregnancies. However, the phenotypic heterogeneity observed in Type 2 GD patients makes the diagnosis a challenge for physicians unfamiliar with the disease. In general, Type 2 GD should be considered in the differential diagnosis of infants presenting with splenomegaly, failure to thrive, congenital ichthyosis, hydrops fetalis and in families with recurrent fetal losses. Reliable methods of diagnosing GD include measurement of glucocerebrosidase enzymatic activity and direct sequencing of *GBA1*. Confirmation of the diagnosis of Type 2 GD will direct treatment considerations and allow physicians to focus on critical clinical issues. Treating these infants is a challenge, and it is important to involve the medical team, family, and support networks early on, in order to identify goals of treatment and help families to cope with this grave diagnosis.

Acknowledgments

This work was supported by the Intramural Research Programs of the National Human Genome Research Institute, and the National Institutes of Health. The authors also greatly acknowledge the many patients, family members and referring physicians whose generous contributions enabled us to compose this work.

References

- [1] B.E. Rosenbloom, N.J. Weinreb, Gaucher disease: a comprehensive review, *Crit. Rev. Oncog.* 18 (2013) 163–175.
- [2] E. Sidransky, Gaucher disease: complexity in a "simple" disorder, *Mol. Genet. Metab.* 83 (2004) 6–15, <http://dx.doi.org/10.1016/j.ymgme.2004.08.015>.
- [3] E. Sidransky, Gaucher disease: insights from a rare mendelian disorder, *Discov. Med.* 14 (2012) 273–281.
- [4] D.L. Stone, E. Sidransky, Hydrops fetalis: lysosomal storage disorders in extremis, *Adv. Pediatr.* 46 (1999) 409–440.
- [5] N. Tayebi, D.L. Stone, E. Sidransky, Type 2 Gaucher disease: an expanding phenotype, *Mol. Genet. Metab.* 68 (1999) 209–219, <http://dx.doi.org/10.1006/mgme.1999.2918>.
- [6] N. Gupta, I.M. Oppenheim, E.F. Kauvar, N. Tayebi, E. Sidransky, Type 2 Gaucher disease: phenotypic variation and genotypic heterogeneity, *Blood Cells Mol. Dis.* 46 (2011) 75–84, <http://dx.doi.org/10.1016/j.bcmd.2010.08.012>.
- [7] C. Mignot, A. Gelot, B. Bessières, F. Daffos, M. Voyer, F. Menez, et al., Perinatal-lethal Gaucher disease, *Am. J. Med. Genet. A* 120A (2003) 338–344, <http://dx.doi.org/10.1002/ajmg.a.20117>.
- [8] N. Tayebi, K.J. Reissner, E.K. Lau, B.K. Stubblefield, A.C. Klineburgess, B.M. Martin, et al., Genotypic heterogeneity and phenotypic variation among patients with type 2 Gaucher's disease, *Pediatr. Res.* 43 (1998) 571–578, <http://dx.doi.org/10.1203/00006450-199805000-00003>.
- [9] E. Sidransky, M. Fartasch, R.E. Lee, L.A. Metlay, S. Abella, A. Zimran, et al., Epidermal abnormalities may distinguish Type 2 from Type 1 and Type 3 of Gaucher disease, *Pediatr. Res.* 39 (1996) 134–141, <http://dx.doi.org/10.1203/00006450-199601000-00020>.
- [10] W.M. Holleran, E.I. Ginns, G.K. Menon, J.-U. Grundmann, M. Fartasch, C.E. McKinney, et al., Consequences of B-glucocerebrosidase deficiency in epidermis, *J. Clin. Invest.* 93 (1994) 1756–1764.
- [11] K.S. Hruska, M.E. LaMarca, C.R. Scott, E. Sidransky, Gaucher disease: mutation and polymorphism spectrum in the glucocerebrosidase gene (*GBA*), *Hum. Mutat.* 29 (2008) 567–583, <http://dx.doi.org/10.1002/humu.20676>.
- [12] N.L.S. Tang, W. Zhang, G.A. Grabowski, K.F. To, F.Y.M. Choy, S.L. Ma, et al., Novel mutations in type 2 Gaucher disease in Chinese and their functional characterization by heterologous expression, *Hum. Mutat.* 26 (2005) 59–60, <http://dx.doi.org/10.1002/humu.9348>.
- [13] D.L. Stone, N. Tayebi, E. Orvisky, B. Stubblefield, V. Madike, E. Sidransky, Glucocerebrosidase gene mutations in patients with type 2 Gaucher disease, *Hum. Mutat.* 15 (2000) 181–188.
- [14] S.A. Scott, L. Edelmann, L. Liu, M. Luo, R.J. Desnick, R. Kornreich, Experience with carrier screening and prenatal diagnosis for 16 Ashkenazi Jewish genetic diseases, *Hum. Mutat.* 31 (2010) 1240–1250, <http://dx.doi.org/10.1002/humu.21327>.
- [15] V. Koprivica, D.L. Stone, J.K. Park, M. Callahan, A. Frisch, I.J. Cohen, et al., Analysis and classification of 304 mutant alleles in patients with type 1 and type 3 Gaucher disease, *Am. J. Hum. Genet.* 66 (2000) 1777–1786, <http://dx.doi.org/10.1086/302925>.
- [16] E. Beutler, H. Liebman, T. Gelbart, E. Stefanski, Three Gaucher-disease-producing mutations in a patient with Gaucher disease: mechanism and diagnostic implications, *Acta Haematol.* 104 (2000) 103–105, <http://dx.doi.org/10.1159/000039760>.
- [17] J. Vithayathil, G. Gibney, A.D. Baxevanis, B.K. Stubblefield, E. Sidransky, N. Tayebi, Glucocerebrosidase mutation H255Q appears to be exclusively in cis with D409H: structural implications, *Clin. Genet.* 75 (2009) 503–504, <http://dx.doi.org/10.1111/j.1399-0004.2009.01163.x>.
- [18] E. Sidransky, M.A. Nalls, J.O. Aasly, J. Aharon-Peretz, G. Annesi, E.R. Barbosa, et al., Multicenter analysis of glucocerebrosidase mutations in Parkinson's disease, *N. Engl. J. Med.* 361 (2009) 1651–1661, <http://dx.doi.org/10.1056/NEJMoa0901281>.
- [19] E. Orvisky, J.K. Park, M.E. LaMarca, E.I. Ginns, B.M. Martin, N. Tayebi, et al., Glucosylsphingosine accumulation in tissues from patients with Gaucher disease: correlation with phenotype and genotype, *762002*. 262–270.
- [20] T. Farfel-Becker, E.B. Vitner, S.N.R. Pressey, R. Eilam, J.D. Cooper, A.H. Futerman, Spatial and temporal correlation between neuron loss and neuroinflammation in a mouse model of neuronopathic Gaucher disease, *202011*. 1375–1386, <http://dx.doi.org/10.1093/hmg/ddr019>.
- [21] T. Farfel-Becker, E.B. Vitner, S.L. Kelly, J.R. Bame, J. Duan, V. Shinder, et al., Neuronal accumulation of glucosylceramide in a mouse model of neuronopathic Gaucher disease leads to neurodegeneration, *Hum. Mol. Genet.* 23 (2014) 843–854, <http://dx.doi.org/10.1093/hmg/ddt468>.
- [22] E.B. Vitner, R. Salomon, T. Farfel-Becker, A. Meshcheriakova, M. Ali, A.D. Klein, et al., RIPK3 as a potential therapeutic target for Gaucher's disease, *Nat. Med.* 20 (2014) 204–208, <http://dx.doi.org/10.1038/nm.3449>.
- [23] W. Zhou, J. Yuan, Necroptosis in health and diseases, *Semin. Cell Dev. Biol.* (2014) 1–10, <http://dx.doi.org/10.1016/j.semcdb.2014.07.013>.
- [24] A. Tyłki-Szymańska, A. Vellodi, A. El-Beshlawy, J.A. Cole, E. Kolodny, Neuronopathic Gaucher disease: demographic and clinical features of 131 patients enrolled in the International Collaborative Gaucher Group Neurological Outcomes Subregistry, *J. Inher. Metab. Dis.* 33 (2010) 339–346, <http://dx.doi.org/10.1007/s10545-009-9009-6>.
- [25] L.A. Newman, C. Keckley, M.C. Petersen, A. Hammer, Swallowing function and medical diagnoses in infants suspected of dysphagia, *Pediatrics* 108 (2001) e106–e106, <http://dx.doi.org/10.1542/peds.108.6.e106>.
- [26] R. Hanna, M.T. McDonald, J.A. Sullivan, J.F. Mackey, V. Krishnamurthy, P.S. Kishnani, Diagnostic and treatment challenges of neuronopathic Gaucher disease: two cases with an intermediate phenotype, *J. Inher. Metab. Dis.* (2004) 687–690.
- [27] J.D. Tobias, R. Atwood, S. Lowe, G.W. Holcomb, Anesthetic considerations in the child with Gaucher disease, *J. Clin. Anesth.* 5 (1993) 150–153.
- [28] C.A. Prows, N. Sanchez, C. Daugherty, G.A. Grabowski, Gaucher disease: enzyme therapy in the acute neuronopathic variant, *Am. J. Med. Genet.* 71 (1997) 16–21,

- [http://dx.doi.org/10.1002/\(SICI\)1096-8628\(19970711\)71:1<16::AID-AJMG3>3.0.CO;2-O](http://dx.doi.org/10.1002/(SICI)1096-8628(19970711)71:1<16::AID-AJMG3>3.0.CO;2-O).
- [29] C. Mignot, D. Doummar, I. Maire, T.B. De Villemeur, Type 2 Gaucher disease: 15 new cases and review of the literature, *282006*. 39–48, <http://dx.doi.org/10.1016/j.braindev.2005.04.005>.
- [30] A. Rapoport, Sublingual atropine drops for the treatment of pediatric sialorrhea, *J. Pain Symptom Manag.* 40 (2010) 783–788, <http://dx.doi.org/10.1016/j.jpainsymman.2010.02.007>.
- [31] A. Erikson, B. Bembi, R. Schiffmann, Neuronopathic forms of Gaucher's disease, *Baillieres Clin. Haematol.* 10 (1997) 711–723.
- [32] C.J. Wusthoff, R.A. Shellhaas, D.J. Licht, Management of common neurologic symptoms in pediatric palliative care: seizures, agitation, and spasticity, *Pediatr. Clin. N. Am.* 54 (2007) 709–733, <http://dx.doi.org/10.1016/j.pcl.2007.06.004>.
- [33] C.-Y. Chung, C.-L. Chen, A.M.-K. Wong, Pharmacotherapy of spasticity in children with cerebral palsy, *J. Formos. Med. Assoc.* 110 (2011) 215–222, [http://dx.doi.org/10.1016/S0929-6646\(11\)60033-8](http://dx.doi.org/10.1016/S0929-6646(11)60033-8).
- [34] J.K. Park, E. Orvisky, N. Tayebi, C. Kaneski, M.E. LaMarca, B.K. Stubblefield, et al., Myoclonic epilepsy in Gaucher disease: genotype–phenotype insights from a rare patient subgroup, *Pediatr. Res.* 53 (2003) 387–395, <http://dx.doi.org/10.1203/01.PDR.0000049515.79882.94>.
- [35] A.T. Berg, C.B. Baca, T. Loddenkemper, D. Dlugos, Priorities in pediatric epilepsy research, *Neurology* (2013) 1166–1175.
- [36] H.C. Glass, Neonatal seizures, *Clin. Perinatol.* 41 (2014) 177–190, <http://dx.doi.org/10.1016/j.clp.2013.10.004>.
- [37] O. Dulac, B. Plecko, S. Gataullina, N.I. Wolf, Occasional seizures, epilepsy, and inborn errors of metabolism, *Lancet Neurol.* 13 (2014) 727–739, [http://dx.doi.org/10.1016/S1474-4422\(14\)70110-3](http://dx.doi.org/10.1016/S1474-4422(14)70110-3).
- [38] S. Szlam, M. Meredith, Shake, rattle, and roll: an update on pediatric seizures, *Pediatr. Emerg. Care* 29 (2013) 1287–1294.
- [39] A.F. Elias, M.R. Johnson, J.K. Boitnott, D. Valle, Neonatal cholestasis as initial manifestation of type 2 Gaucher disease: a continuum in the spectrum of early onset Gaucher disease, *JIMD Rep.* 5 (2011) 95–98, http://dx.doi.org/10.1007/8904_2011_104.
- [40] C. Barbier, L. Devisme, D. Dobbelaere, O. Noizet, B. Nelken, F. Gottrand, Neonatal cholestasis and infantile Gaucher disease: a case report, *Acta Paediatr.* (2002) 1399–1401.
- [41] H. Kelsey, C. Christopoulos, A.A. Gray, S.J. Machin, Acquired pseudo–pseudo Bernard–Soulier syndrome complicating Gaucher's disease, *J. Clin. Pathol.* (1994) 162–165.
- [42] H. CE, M. Levi, F. Berends, J.M. Aerts, M.H. van Oers, Coagulation abnormalities in type 1 Gaucher disease are due to low-grade activation and can be partly restored by enzyme supplementation therapy, *Br. J. Haematol.* (1996) 470–476.
- [43] B. Gulhan, U. Ozcelik, F. Gurakan, S. Gucer, D. Orhan, G. Cinel, et al., Different features of lung involvement in Niemann–Pick disease and Gaucher disease, *Respir. Med.* 106 (2012) 1278–1285, <http://dx.doi.org/10.1016/j.rmed.2012.06.014>.
- [44] A. Erikson, K. Johansson, J.E. Mansson, L. Scennerholm, Enzyme replacement therapy of infantile Gaucher disease, *Neuropediatrics* (1993) 237–238.
- [45] A. Vellodi, A. Tylki-Szymanska, E.H. Davies, E. Kolodny, B. Bembi, T. Collin-Histed, et al., Management of neuronopathic Gaucher disease: revised recommendations, *J. Inher. Metab. Dis.* 32 (2009) 660–664, <http://dx.doi.org/10.1007/s10545-009-1164-2>.
- [46] D. Elstein, A. Abrahamov, A. Zimran, Ethical considerations for enzyme replacement therapy in neuronopathic Gaucher disease, *Clin. Genet.* 54 (1998) 179–184.
- [47] B. Bembi, G. Ciana, M. Zanatta, Cerebrospinal-fluid infusion of algucerase in the treatment for acute neuronopathic Gaucher's disease, *Pediatr. Res.* (1995) A425.
- [48] R.R. Lonser, S. Walbridge, G.J. Murray, M.R. Aizenberg, A.O. Vortmeyer, J.M.F.G. Aerts, et al., Convection perfusion of glucocerebrosidase for neuronopathic Gaucher's disease, *Ann. Neurol.* 57 (2005) 542–548, <http://dx.doi.org/10.1002/ana.20444>.
- [49] R.R. Lonser, R. Schiffman, R.A. Robison, J.A. Butman, Z. Quezado, M.L. Walker, et al., Image-guided, direct convective delivery of glucocerebrosidase for neuronopathic Gaucher disease, *Neurology* 68 (2007) 254–261, <http://dx.doi.org/10.1212/01.wnl.0000247744.10990.e6>.
- [50] J.A. Barranger, S.I. Rapoport, W.R. Fredericks, P.G. Pentchev, K.D. MacDermot, J.K. Steusing, et al., Modification of the blood–brain barrier: increased concentration and fate of enzymes entering the brain, *Proc. Natl. Acad. Sci. U.S.A.* 76 (1979) 481–485.
- [51] R. Schiffmann, E.J. FitzGibbon, C. Harris, C. DeVile, E.H. Davies, L. Abel, et al., Randomized, controlled trial of miglustat in Gaucher's disease type 3, *Ann. Neurol.* 64 (2008) 514–522, <http://dx.doi.org/10.1002/ana.21491>.
- [52] S. Ito, P. Pophali, W. CO, E.K. Koklanaris, J. Superata, G.A. Fahle, et al., CMV reactivation is associated with a lower incidence of relapse after allo-SCT for CML, *Bone Marrow Transplant.* 48 (2013) 1313–1316, <http://dx.doi.org/10.1038/bmt.2013.49>.
- [53] S. Ito, A.J. Barrett, Gauchers disease—a reappraisal of hematopoietic stem cell transplantation, *Pediatr. Hematol. Oncol.* 30 (2013) 61–70, <http://dx.doi.org/10.3109/08880018.2012.762076>.
- [54] O. Ringdén, C.G. Groth, A. Erikson, S. Granqvist, J.E. Mansson, E. Sparrelid, Ten years' experience of bone marrow transplantation for Gaucher disease, *Transplantation* 59 (1995) 864–870.
- [55] M. Machaczka, Allogeneic hematopoietic stem cell transplantation for treatment of Gaucher disease, *Pediatr. Hematol. Oncol.* 30 (2013) 459–461, <http://dx.doi.org/10.3109/08880018.2013.793757>.
- [56] H. Saranjam, S.S. Chopra, H. Levy, B. Stubblefield, E. Maniawang, I.J. Cohen, et al., A germline or de novo mutation in two families with Gaucher disease: implications for recessive disorders, *Eur. J. Hum. Genet.* 21 (2013) 115–117, <http://dx.doi.org/10.1038/ejhg.2012.105>.
- [57] W.S. Benko, K.S. Hruska, N. Nagan, O. Goker-Alpan, P.S. Hart, R. Schiffmann, et al., Uniparental disomy of chromosome 1 causing concurrent Charcot–Marie–Tooth and Gaucher disease type 3, *Neurology* 70 (2008) 976–978, <http://dx.doi.org/10.1212/01.wnl.0000305963.37449.32>.
- [58] M.A. Cabrera-Salazar, M. DeRiso, S.D. Bercury, L. Li, J.T. Lydon, W. Weber, et al., Systemic delivery of a glucosylceramide synthase inhibitor reduces CNS substrates and increases lifespan in a mouse model of type 2 Gaucher disease, *PLoS ONE* 7 (2012) e43310, <http://dx.doi.org/10.1371/journal.pone.0043310>.
- [59] R.Y. Wang, O.A. Bodamer, M.S. Watson, W.R. Wilcox, Lysosomal storage diseases: diagnostic confirmation and management of presymptomatic individuals, *Genet. Med.* 13 (2011) 457–484, <http://dx.doi.org/10.1097/GIM.0b013e318211a7e1>.
- [60] Y. Suzuki, Chaperone therapy update: Fabry disease, GM1-gangliosidosis and Gaucher disease, *Brain Dev.* 35 (2013) 515–523, <http://dx.doi.org/10.1016/j.braindev.2012.12.002>.
- [61] R. Giugliani, S. Waldek, D.P. Germain, K. Nicholls, D.G. Bichet, J.K. Simosky, et al., A Phase 2 study of migalastat hydrochloride in females with Fabry disease: selection of population, safety and pharmacodynamic effects, *Mol. Genet. Metab.* 109 (2013) 86–92, <http://dx.doi.org/10.1016/j.ymgme.2013.01.009>.
- [62] A. Joosten, C. Decroocq, J. de Sousa, J.P. Schneider, E. Etame, A. Bodlenner, et al., A systematic investigation of lminosugar click clusters as pharmacological chaperones for the treatment of Gaucher disease, *ChemBioChem* 15 (2014) 309–319, <http://dx.doi.org/10.1002/cbic.201300442>.
- [63] W. Schonemann, E. Gallienne, K. Ikeda-Obatake, N. Asano, S. Nakagawa, A. Kato, et al., Glucosylceramide mimics: highly potent GCse inhibitors and selective pharmacological chaperones for mutations associated with types 1 and 2 Gaucher disease, *ChemMedChem* 8 (2013) 1805–1817, <http://dx.doi.org/10.1002/cmdc.201300327>.
- [64] G. Tiscornia, E. Lorenzo Vivas, L. Matalonga, I. Berniakovich, M. Barragan Monasterio, C. Eguizabal, et al., Neuronopathic Gaucher's disease: induced pluripotent stem cells for disease modelling and testing chaperone activity of small compounds, *Hum. Mol. Genet.* 22 (2013) 633–645, <http://dx.doi.org/10.1093/hmg/ddt471>.
- [65] E. Aflaki, B.K. Stubblefield, E. Maniawang, G. Lopez, N. Moaven, E. Goldin, et al., Macrophage models of Gaucher disease for evaluating disease pathogenesis and candidate drugs, *Sci. Transl. Med.* 6 (2014), <http://dx.doi.org/10.1126/scitranslmed.3008659> (240ra73–240ra73).
- [66] G. Parenti, C. Pignata, P. Vajro, M.C. Salerno, New strategies for the treatment of lysosomal storage diseases (review), *Int. J. Mol. Med.* 31 (2012) 11–20, <http://dx.doi.org/10.3892/ijmm.2012.1187>.
- [67] D.L. Hill, V. Miller, J.K. Walter, K.W. Carroll, W.E. Morrison, D.A. Munson, et al., Regoaling: a conceptual model of how parents of children with serious illness change medical care goals, *BMC Palliat. Care* 13 (2014) 1–8, <http://dx.doi.org/10.1186/1472-684X-13-9>.
- [68] J.M. Youngblut, D. Brooten, Perinatal and pediatric issues in palliative and end-of-life care from the 2011 Summit on the Science of Compassion, *Nurs. Outlook* 60 (2012) 343–350, <http://dx.doi.org/10.1016/j.outlook.2012.08.007>.
- [69] M.E. Macdonald, S. Liben, F.A. Carnevale, J.E. Rennick, S.L. Wolf, D. Meloche, et al., Parental perspectives on hospital staff members "acts of kindness and commemoration after a child's death, *Pediatrics* 116 (2005) 884–890, <http://dx.doi.org/10.1542/peds.2004-1980>.
- [70] K. Lui, C. Commens, R. Choong, R. Jaworski, Colloidion babies with Gaucher's disease, *Arch. Dis. Child.* 63 (1988) 854–856.
- [71] D.L. Stone, W.F. Carey, J. Christodoulou, D. Silence, P. Nelson, M. Callahan, et al., Type 2 Gaucher disease: the colloidion baby phenotype revisited, *Arch. Dis. Child. Fetal Neonatal Ed.* 82 (2000) F163–F166.
- [72] D.L. Stone, O.P. van Diggelen, J.B. de Klerk, J.L. Gaillard, M.F. Niermeijer, R. Willemsen, et al., Is the perinatal lethal form of Gaucher disease more common than classic type 2 Gaucher disease? 71999. 505–509, <http://dx.doi.org/10.1038/sj.ejhg.5200315>.
- [73] S. Haverkaemper, T. Marquardt, I. Hausser, K. Timme, T. Kuehn, C. Hertzberg, et al., Congenital ichthyosis in severe type II Gaucher disease with a homozygous null mutation, *Neonatology* 100 (2011) 194–197, <http://dx.doi.org/10.1159/000324116>.
- [74] A. Chan, W.M. Holleran, T. Ferguson, D. Crumrine, O. Goker-Alpan, R. Schiffmann, et al., Skin ultrastructural findings in type 2 Gaucher disease: diagnostic implications, *Mol. Genet. Metab.* 104 (2011) 631–636, <http://dx.doi.org/10.1016/j.ymgme.2011.09.008>.
- [75] K. Reissner, N. Tayebi, B.K. Stubblefield, V. Koprivica, M. Blitzer, W. Holleran, et al., Type 2 Gaucher disease with hydrops fetalis in an Ashkenazi Jewish family resulting from a novel recombinant allele and a rare splice junction mutation in the glucocerebrosidase locus, *Mol. Genet. Metab.* 63 (1998) 281–288, <http://dx.doi.org/10.1006/mgme.1998.2675>.
- [76] H.J. Church, A. Cooper, F. Stewart, C.M. Thornton, J.E. Wraith, Homozygous loss of a cysteine residue in the glucocerebrosidase gene results in Gaucher's disease with a hydroptic phenotype, *Eur. J. Hum. Genet.* 12 (2004) 975–978, <http://dx.doi.org/10.1038/sj.ejhg.5201251>.
- [77] H. Zhao, M. Keddache, L. Bailey, G. Arnold, G. Grabowski, Gaucher's disease: identification of novel mutant alleles and genotype–phenotype relationships, *Clin. Genet.* 64 (2003) 57–64.
- [78] H. Michelakakis, E. Dimitriou, M. Moraitou, M. Valari, E. Yatrakou, V. Mitsiadi, et al., Perinatal lethal form of Gaucher disease. Clinical and molecular characterization of a Greek case, *Blood Cells Mol. Dis.* 44 (2010) 82–83, <http://dx.doi.org/10.1016/j.bcmd.2009.11.007>.
- [79] A. Chabás, L. Gort, A. Díaz-Font, M. Montfort, R. Santamaría, M. Cidrás, et al., Perinatal lethal phenotype with generalized ichthyosis in a type 2 Gaucher disease patient with the [L444P;E326K]/P182L genotype: effect of the E326K change in neonatal and classic forms of the disease, *Blood Cells Mol. Dis.* 35 (2005) 253–258, <http://dx.doi.org/10.1016/j.bcmd.2005.04.007>.

- [80] G. Sinclair, F.Y. Choy, L. Humphries, A novel complex allele and two new point mutations in type 2 (acute neuronopathic) Gaucher disease, *Blood Cells Mol. Dis.* 24 (1998) 420–427, <http://dx.doi.org/10.1006/bcmd.1998.0210>.
- [81] O. Goker-Alpan, R. Schiffmann, J.K. Park, B.K. Stubblefield, N. Tayebi, E. Sidransky, Phenotypic continuum in neuronopathic Gaucher disease: an intermediate phenotype between type 2 and type 3, *J. Pediatr.* 143 (2003) 273–276, [http://dx.doi.org/10.1067/S0022-3476\(03\)00302-0](http://dx.doi.org/10.1067/S0022-3476(03)00302-0).
- [82] S. Aviner, B.-Z. Garty, A. Rachmel, H.N. Baris, E. Sidransky, A. Shuffer, et al., Type 2 Gaucher disease occurs in Ashkenazi Jews but is surprisingly rare, *Blood Cells Mol. Dis.* 43 (2009) 294–297, <http://dx.doi.org/10.1016/j.bcmd.2009.08.004>.
- [83] S. Rowlands, H. Murray, Prenatal ultrasound findings in a fetus diagnosed with Gaucher's disease (type 2) at birth, *Prenat. Diagn.* 17 (1997) 765–769, [http://dx.doi.org/10.1002/\(SICI\)1097-0223\(199708\)17:8<765::AID-PD122>3.0.CO;2-3](http://dx.doi.org/10.1002/(SICI)1097-0223(199708)17:8<765::AID-PD122>3.0.CO;2-3).
- [84] E. Sidransky, N. Tayebi, B.K. Stubblefield, W. Eliason, A. Klineburgess, G.P. Pizzolato, et al., The clinical, molecular, and pathological characterisation of a family with two cases of lethal perinatal type 2 Gaucher disease, *J. Med. Genet.* 33 (1996) 132–136.
- [85] E. Sidransky, D.M. Sherer, E.I. Ginns, Gaucher Disease in the neonate: a distinct Gaucher phenotype is analogous to a mouse model created by targeted disruption of the glucocerebrosidase gene, *Pediatr. Res.* 32 (1992) 1–5.